

ZONING AND LAND USE

870 Attachment 1

City of Camden

Appendix I Invasive Nonindigenous Plant Species

The following trees, shrubs, vines and herbaceous plants as well as their cultivars are known or believed to be invasive to natural ecosystems in New Jersey. With the exception of common reed (*Phragmites australis*), which is cosmopolitan but may include nonindigenous lineages known to aggressively invade and dominate diverse wetland habitats, all species listed are nonindigenous. While many of these species may be available at commercial nurseries for landscaping, because of their invasive nature, they are not suitable for landscaping, replanting of facilities, or ecological restorations. For each species the common name and scientific name are provided. Scientific names should be consulted for accuracy in identifying species. Note that several species may be included under a single common name. The list will be periodically updated based on new information.

Trees	
Black locust	(<i>Robinia pseudoacacia</i>)
Callery Pear	(<i>Pyrus calleryana</i>)
Catalpa	(<i>Catalpa bignonioides</i>)
Chinese	(<i>Ulmus parvifolia</i>)
Crack willow	(<i>Salix fragilis</i>)
Devil's walking stick	(<i>Aralia spinosa</i>)
Empress tree	(<i>Paulownia tomentosa</i>)
Japanese cork tree	(<i>Phellodendron japonicum</i>)
Mimosa	(<i>Albizia julibrissin</i>)
Norway maple	(<i>Acer platanoides</i>)
Paper-mulberry	(<i>Broussonetia papyrifera</i>)
Scotch pine	(<i>Pinus sylvestris</i>)
Siberian elm	(<i>Ulmus pumila</i>)
Sweet cherry	(<i>Prunus avium</i>)
Tree of heaven	(<i>Ailanthus altissima</i>)
Umbrella tree	(<i>Magnolia tripetala</i>)
White mulberry	(<i>Morus alba</i>)
White poplar	(<i>Populus alba</i>)
White willow	(<i>Salix alba</i>)
Yellow buckeye	(<i>Aesculus flava</i>)

CAMDEN CODE

Shrubs	
Amur honeysuckle	(Lonicera maackii)
Autumn olive	(Elaeagnus umbrellata)
Black alder	(Alnus glutinosa)
Bramble	(Rubus discolor)
Bush honeysuckles	(Lonicera morrowii)
Butterfly-bush	(Buddleja davidii)
Clammy locust	(Robibia viscosa)
Common buckthorn	(Rhamnus cathartica)
Coral-berry	(Symphoricarpos orbiculatus)
Dog rose	(Rosa canina)
European barberry	(Berberis vulgaris)
European spindle-tree	(Euonymus europaeus)
False indigo-bush	(Amorpha fruticosa)
Fragrant sumac	(Rhus aromatica)
Garden red currant	(Ribes sativum)
Japanese barberry	(Berberis thunbergii)
Japanese holly	(Ilex crenata)
Jethead	(Rhodotypos scandens)
Linden arrow-wood	(Viburnum dilatatum)
Morrow's bush honeysuckle	(Lonicera morrowii)
Multiflora rose	(Rosa multiflora)
Oriental redbud	(Photinia villosa)
Privet	(Ligustrum obtusifolium)
Rose acacia	(Robinia hispida)
Rugosa rose	(Rosa rugosa)
Russian olive	(Elaeagnus angustifolia)
Scotch broom	(Cytisus scoparius)
Shrub lespedeza	(Lespedeza thunbergii)
Siebold viburnum	(Viburnum sieboldii)
Smooth buckthorn	(Rhamnus frangula)
Snowberry	(Symphoricarpos albus var. laevigatus)
Sweetbrier	(Rosa micrantha)
Tartarian honeysuckle	(Lonicera tatarica)
Winged burning bush	(Euonymus alatus)
Winter creeper	(Euonymus fortunei)

ZONING AND LAND USE

Vines	
Akebia	(Akebia quinata)
Black swallow-wort	(Cynanchum louiseae)
English ivy	(Hedera helix)
Japanese honeysuckle	(Lonicera japonica var. chinensis)
Kudzu	(Pueraria lobata)
Mile-a-minute	(Polygonum perfoliatum)
Matrimony vine	(Lycium barbarum)
Oriental bittersweet	(Celastrus orbiculatus)
Periwinkle	(Vinca minor)
Porcelain-berry	(Ampelopsis brevipedunculata)
Wisteria	(Wisteria floribunda)
Yam-leaved clematis	(Clematis terniflora)

Herbs	
Bedstraw	(Galium mollugo)
Bitter-cress	(Cardamine impatiens)
Black-eyed Susan	(Rudbeckia hirta var. pulcherrima)
Bull thistle	(Cirsium vulgare)
Bush clover	(Lespedeza cuneata)
Butter and eggs	(Linaria vulgaris)
Canada thistle	(Cirsium arvense)
Chickweed	(Cerastium bieberstenii)
Chicory	(Cichorium intybus)
Coltsfoot	(Tussilago farfara)
Cottonweed	(Froelichia gracilis)
Creeping primrose-willow	(Ludwigia peploides var. glabrescens)
Crown-vetch	(Coronilla varia)
Curly dock	(Rumex crispus)
Cypress spurge	(Euphorbia cyparissias)
Dame' rocket	(Hesperis matronalis)
Dusty miller	(Artemisia stelleriana)
Eurasian water-milfoil	(Myriophyllum spicatum)
Field garlic	(Allium vineale)
Garlic mustard	(Alliaria petiolata)
Giant knotweed	(Polygonum sacalinense)
Goosefoot	(Chenopodium ambrosioides)
Goutweed	(Aegopodium podagraria)

CAMDEN CODE

Herbs	
Ground-ivy	(Glechoma hederaceae)
Hawkweed	(Hieracium caespitosum)
Indian strawberry	(Duchesnea indica)
Japanese hops	(Humulus japonicus)
Japanese knotwood	(Polygonum cuspidatum)
Leafy spurge	(Euphorbia escula)
Lesser celandine	(Ranunculus ficaria)
Lobelia	(Lobelia chinensis)
Moneywort	(Lysimachia nummularia)
Mullein	(Verbascum blattaria)
Mugwort	(Artemisia vulgaris)
Orange daylily	(Hemerocallis fulva)
Ox-eye daisy	(Leucantemum vulgare)
Bearded beggarticks	(Bidens aristosa)
Pondweed	(Potamogeton crispus)
Prince's feather	(Polygonum orientale)
Purple loosestrife	(Lythrum salicaria)
Queen Ann's-lace	(Daucus carota)
Ragged-robin	(Lychnis flos-cuculi)
Skeleton-weed	(Chondrilla juncea)
Speedwell	(Veronica beccabunga)
Spotted knapweed	(Centaurea maculosa)
Sweet clover	(Melilotus officinalis)
Teasel	(Dipsacus fullonum)
Thistle	(Carduus acanthoides)
Water-chestnut	(Trapa natans)
Water-starwort	(Callitriche stagnalis)
Water-thyme	(Hydrilla verticillata)
White mullein	(Verbascum lychnitis)
Wild chervil	(Anthriscus sylvestris)
Winter-cress	(Barbarea verna)

Graminoid	
African weeping love grass	(Eragrostis curvula)
Brome	(Bromus japonicus)
Cane	(Arundinaria gigantean)

ZONING AND LAND USE

Graminoid	
Canary grass	(Phalaris canariensis)
Chinese silver grass	(Miscanthus sinensis)
Common reed	(Phragmites australis)
Japanese sedge	(Carex kobomugi)
Japanese stiltgrass	(Microstegium vimineum)
Meadow fescue	(Festuca pratensis)
Small carp grass	(Arthraxon hispidus)
Sweet vernal grass	(Anthoxanthum odoratum)
Umbrella-sedge	(Cyperus amuricus)